

Reply to Comment on “Quantum Phase Transition of Randomly-Diluted Heisenberg Antiferromagnet on a Square Lattice”

Syngé Todo^{1*}, Hajime Takayama¹, and Naoki Kawashima²

¹*Institute for Solid State Physics, University of Tokyo, Kashiwa 277-8581, Japan*

²*Department of Physics, Tokyo Metropolitan University, Tokyo 192-0397, Japan*
(February 1, 2008)

In Ref. [1], we have studied the quantum phase transition of the two-dimensional diluted Heisenberg antiferromagnet by using the quantum Monte Carlo method. The two main conclusions in the article are i) the critical concentration of magnetic sites is equal to the classical percolation threshold ($p_{cl} = 0.592746$), and ii) the value of the critical exponents, such as β and Ψ , significantly depends on the spin size S . In [2], Sandvik has presented the result of his quantum Monte Carlo simulation for the same system only in the case of $S = \frac{1}{2}$, in which the temperature dependence of the static structure factor is studied quite precisely. Based on his data, he claimed that the data in Ref. [1] should be seriously affected by the finiteness of temperature in our simulation. In addition, he calculated the cluster magnetization $m_{cl}(L)$ just at $p = p_{cl}$, and claimed that it could remain finite even in the thermodynamic limit in the same way as the classical ($S = \infty$) case, although quite large finite-size corrections to its asymptotic behavior should be considered in the quantum case. As we will discuss below, however, his data do not conflict with our data nor with our conclusions. In particular, the data presented in Fig. 1(b) in [2] can be explained from our view point assuming non-magnetic ground states.

As for his first claim on the effect of the temperature, we agree that there may be a small underestimation for the zero-temperature values of the static structure factor. However, even if we replace our data point in Fig. 3 in [1] for $L = 48$ by what we can expect from Fig. 1(a) in [2], it would result in only a minor change in the estimate of Ψ . Therefore, the error caused by this underestimation, if any, does not seem to be large enough to invalidate our conclusions.

In addition, in Ref. [1], we have presented the result of another finite-size scaling analysis (Fig. 4), in which the data at higher temperatures are also used together with the one at the lowest temperature in the simulation. In this analysis, we have explicitly taken into account the finite-temperature effect on the static structure factor by introducing the dynamical exponent z . For $S = 1$, the value of the exponent Ψ coincides with that obtained by the former analysis. On the other hand, for $S = \frac{1}{2}$, it becomes slightly larger (about 9%) than the other estimate based on Fig. 3 in [1]. This small difference may be due to the finite-temperature effect pointed out in [2], though we have not commented on it in [1]. However, even the estimate based on the finite-temperature analysis signif-




FIG. 1. Log-log plot of cluster magnetization (Ref. [2]). The solid line is obtained by least-squares fitting for $L \geq 28$.

icantly differs from those for $S \geq 1$. Therefore, we think our second conclusion on the non-universal behavior can not be excluded.

It was also suggested [2], based on the calculation of the magnetization of the largest connected cluster for each sample, that all the critical exponents, including Ψ and β , are equal to the classical values. The data were plotted against $x \equiv 1/L^{D/2}$, and appeared to suggest a finite value in the thermodynamic limit. From our point of view, however, his data can be interpreted as follows. The quantity $m_{cl}^2(L)$ defined in Ref. [2] should be asymptotically scaled as

$$m_{cl}^2(L) \sim S_s(L, 0, p_{cl})/L^{2D-d}, \quad (1)$$

where $S_s(L, T, p)$ is the static structure factor (Eq. (4) in Ref. [1]) and L^D is the average number of spins in the largest connected clusters in a system of linear size L . According to our results [1], on the other hand, $S_s(L, 0, p_{cl})$ diverges as $L^{2D-d-\alpha}$ with $\alpha = 0.52$ for $S = \frac{1}{2}$, and consequently $m_{cl}^2(L)$ should be scaled as $L^{-0.52}$. In Fig. 1, we show a log-log plot of the cluster magnetization data by Sandvik [2]. One sees that the data for $L \geq 28$ clearly scales as $L^{-0.53}$, which is fully consistent with our results mentioned above.

The authors are grateful to A. W. Sandvik for allowing us to use his numerical data.

* Present address: Theoretical Physics, ETH Zürich, 8093 Zürich, Switzerland. E-mail: wistaria@itp.phys.ethz.ch

[1] K. Kato, S. Todo, K. Harada, N. Kawashima, S. Miyashita, and H. Takayama, Phys. Rev. Lett. **84**, 4204 (2000).

[2] A. W. Sandvik, preprint (cond-mat/0010433).